

DOCTOR DISCUSSION GUIDE FOR SCEMBLIX

Whether you're on treatment or just about to begin, use this discussion guide to start a conversation with your doctor about your medication and find out if SCEMBLIX may be right for you.

1. What questions do you have for your doctor about how to manage your Ph+ CML in chronic phase?

2. What questions do you have for your doctor about setting treatment goals?

3. What questions or concerns do you have about managing your condition with your current treatment or with a treatment like SCEMBLIX?



ASK YOUR DOCTOR IF SCEMBLIX MAY BE RIGHT FOR YOU.

APPROVED USES

SCEMBLIX is a prescription medicine used to treat adults with:

- Newly diagnosed or previously treated Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP)
 - The approval of SCEMBLIX in patients newly diagnosed is based on a study that measured major molecular response (MMR) rate. There are ongoing studies to confirm the clinical benefit of SCEMBLIX for this use and how SCEMBLIX works over a longer period of time

It is not known if SCEMBLIX is safe and effective in children.

IMPORTANT SAFETY INFORMATION

Before taking SCEMBLIX, tell your doctor about all your medical conditions, including if you:

- have a history of inflammation of your pancreas (pancreatitis)
- have a history of heart problems or blood clots in your arteries and veins (types of blood vessels)

Please see Important Safety Information throughout and [click here](#) for accompanying full Prescribing Information, including Patient Information.

 **SCEMBLIX**[®]
(asciminib) 20 mg, 40 mg tablets

IF YOU ARE NEWLY DIAGNOSED WITH Ph+ CML IN CHRONIC PHASE, PLEASE SKIP QUESTIONS 4-6.

4. Which, if any, of the following common side effects from your medication for Ph+ CML in chronic phase are impacting your daily life? (Check all that apply.)

- Fever
- Diarrhea
- Nausea/vomiting
- Bone, muscle, and joint pain
- Difficulty breathing/shortness of breath
- Fluid retention/swelling
- Fatigue that disrupts daily activities
- Unusual bleeding/bruising
- Increased blood pressure
- Rash/itching
- Headache
- Brain fog/confusion
- None

Not on the list?

Add any additional side effects you’re experiencing below and share with your doctor:

IMPORTANT SAFETY INFORMATION (continued)

Before taking SCEMBLIX, tell your doctor about all your medical conditions, including if you: (continued)

- are pregnant or plan to become pregnant. SCEMBLIX can harm your unborn baby

Females who are able to become pregnant:

- Your doctor will do a pregnancy test before you start treatment with SCEMBLIX
- You should use effective birth control (contraception) during treatment and for 1 week after your last dose of SCEMBLIX. Talk to your doctor about birth control methods that may be right for you
- Tell your doctor right away if you become pregnant or think you may be pregnant during treatment with SCEMBLIX
- are breastfeeding or plan to breastfeed. It is not known if SCEMBLIX passes into your breast milk. Do not breastfeed during treatment and for 1 week after your last dose of SCEMBLIX

Tell your doctor about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements.

Taking SCEMBLIX with certain other medicines may affect each other causing side effects and may affect the way that SCEMBLIX works.

Please see Important Safety Information throughout and [click here](#) for accompanying full Prescribing Information, including Patient Information.



5. How frequently do you experience side effects from your medication, if at all?

6. How do the side effects from your medication interfere with your daily life?

Provide your doctor with specific examples:

TO SEE IF SCEMBLIX COULD BE RIGHT FOR YOU, VISIT SCEMBLIX.COM

IMPORTANT SAFETY INFORMATION (continued)

SCEMBLIX may cause serious side effects, including:

- **Low blood cell counts:** SCEMBLIX may cause low platelet counts (thrombocytopenia), low white blood cell counts (neutropenia), and low red blood cell counts (anemia). Your doctor will do blood tests to check your blood cell counts every 2 weeks for the first 3 months of treatment and then monthly or as needed during treatment with SCEMBLIX. Tell your doctor right away if you have unexpected bleeding or easy bruising, blood in your urine or stools, fever, or any signs of an infection
- **Pancreas problems:** SCEMBLIX may increase enzymes in your blood called amylase and lipase, which may be a sign of inflammation of the pancreas (pancreatitis). Your doctor may do blood tests monthly or as needed during treatment with SCEMBLIX to check for problems with your pancreas. Tell your doctor right away if you have severe stomach-area (abdominal) pain or discomfort, nausea, or vomiting
- **High blood pressure:** Your doctor may check your blood pressure and treat any high blood pressure during treatment with SCEMBLIX as needed. Tell your doctor if you develop high blood pressure or symptoms of high blood pressure including confusion, headaches, dizziness, chest pain, or shortness of breath
- **Allergic reaction:** Stop taking SCEMBLIX and get medical help right away if you get any signs or symptoms of an allergic reaction, including:
 - trouble breathing or swallowing
 - swelling of the face, lips, or tongue
 - skin rash or flushing of your skin
 - feeling dizzy or faint
 - fever
 - fast heartbeat

Please see Important Safety Information throughout and [click here](#) for accompanying full Prescribing Information, including Patient Information.



IMPORTANT SAFETY INFORMATION (continued)

SCEMBLIX may cause serious side effects, including: (continued)

• **Heart and blood vessel (cardiovascular) problems:** SCEMBLIX may cause heart and blood vessel problems, including heart attack, stroke, blood clots or blockage of your arteries, heart failure, and abnormal heartbeat, which can be serious and may sometimes lead to death. In the majority of cases, these heart and blood vessel problems happened in people with risk factors or a history of these problems, and previously treated with other tyrosine kinase inhibitor (TKI) medicines. Your doctor may monitor you for heart and blood vessel problems and treat you as needed during treatment with SCEMBLIX. Tell your doctor or get medical help right away if you get:

- shortness of breath
- chest pain or pressure
- a feeling like your heart is beating too fast or you feel abnormal heartbeats
- swelling in your ankles or feet
- dizziness
- weight gain
- numbness or weakness on one side of your body
- decreased vision or loss of vision
- trouble talking
- pain in your arms, legs, back, neck, or jaw
- headache
- severe stomach-area (abdominal) pain

The most common side effects of SCEMBLIX include:

- muscle, bone, or joint pain
- rash
- tiredness
- nose, throat, or sinus (upper respiratory tract) infections
- headache
- stomach-area (abdominal) pain
- diarrhea
- decreased white blood cell counts, platelet counts, and red blood cell counts
- decreased blood calcium corrected levels
- increased blood pancreas enzyme (lipase and amylase) levels
- increased blood fat (cholesterol and triglycerides) levels
- increased blood uric acid levels
- increased blood liver enzyme levels
- increased blood alkaline phosphatase levels
- increased blood creatine kinase levels

Your doctor may change your dose, temporarily stop, or permanently stop treatment with SCEMBLIX if you have certain side effects.

SCEMBLIX may cause fertility problems in females. This may affect your ability to have a child. Talk to your doctor if this is a concern for you.

These are not all the possible side effects of SCEMBLIX. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please [click here](#) for accompanying full Prescribing Information, including Patient Information.

